



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011

Prevalence of age-related hearing loss in Europe: a review

Roth, T N ; Hanebuth, D ; Probst, R

Abstract: Populations are becoming progressively older thus presenting symptoms of diminished organ function due to degenerative processes. These may be physiological or caused by additional factors damaging the organ. Presbycusis refers to the physiological age-related changes of the peripheral and central auditory system leading to hearing impairment and difficulty understanding spoken language. In contrast to epidemiological data of other continents, the prevalence of age-related hearing loss (ARHL) in Europe is not well defined, due in part to the use of different classification systems. We performed a systematic literature review with the aim of gaining a picture of the prevalence of ARHL in Europe. The review included only population and epidemiological studies in English since 1970 with samples in European countries with subjects aged 60 years and above. Nineteen studies met our selection criteria and additional five studies reported self-reported hearing impairment. When these data were crudely averaged and interpolated, roughly 30% of men and 20% of women in Europe were found to have a hearing loss of 30 dB HL or more by age 70 years, and 55% of men and 45% of women by age 80 years. Apparent problems in comparing the available data were the heterogeneity of measures and cut-offs for grades of hearing impairment. Our systematic review of epidemiological data revealed more information gaps than information that would allow gaining a meaningful picture of prevalence of ARHL. The need for standardized procedures when collecting and reporting epidemiological data on hearing loss has become evident. Development of hearing loss over time in conjunction with the increase in life expectancy is a major factor determining strategies of detection and correction of ARHL. Thus, we recommend using the WHO classification of hearing loss strictly and including standard audiometric measures in population-based health surveys.

DOI: <https://doi.org/10.1007/s00405-011-1597-8>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-48826>

Journal Article

Accepted Version

Originally published at:

Roth, T N; Hanebuth, D; Probst, R (2011). Prevalence of age-related hearing loss in Europe: a review. *European Archives of Oto-Rhino-Laryngology*, 268(8):1101-1107.

DOI: <https://doi.org/10.1007/s00405-011-1597-8>

Manuscript Number:

Title: Prevalence of age-related hearing loss (ARHL) in Europe - A Review

Article Type: Review Article (Original Article)

Keywords: Presbycusis, age-related hearing loss, auditory system, cochlear function, epidemiology, Europe, AHEAD

Corresponding Author: Thomas N Roth, MD

Corresponding Author's Institution:

First Author: Thomas N Roth, MD

Order of Authors: Thomas N Roth, MD;Dirk Hanebuth, PhD;Rudolf Probst, MD, Professor

Abstract: Hearing loss is the most widespread sensory impairment in aging people, but population-based epidemiological knowledge or trends of the amount of HL are not well described. A systematic literature review was carried out with the aim to gain a picture of the epidemiology of age related hearing loss in Europe. We searched for prevalence data defining hearing loss by an audiometric threshold between 25 and 45 dB HL for the better ear or self-reported hearing impairment. Population based studies in English reported since 1970 with subjects aged 60 years and above were considered. 19 studies reporting audiometrically measured hearing loss and 5 studies reporting self-reported hearing impairment met our selection criteria. They did not allow establishing an integrative quantitative overview of prevalence rates over age and hearing loss for Europe, but they reflected the well-known patterns of a nonlinear increase of hearing loss with age affecting men more than women. Roughly 30% of men and 20% of women in Europe have a hearing loss of 30 dB HL or more at age 70 years, and 55% of men and 45% of women at age 80 years. The search confirms the known fact that ARHL is a major health concern in the aging population of Europe, but important information gaps became evident. They demonstrate the need of systematic and standardized collections of epidemiological data on hearing loss.

Abstract

Hearing loss is the most widespread sensory impairment in aging people, but population-based epidemiological knowledge or trends of the amount of HL are not well described. A systematic literature review was carried out with the aim to gain a picture of the epidemiology of age related hearing loss in Europe. We searched for prevalence data defining hearing loss by an audiometric threshold between 25 and 45 dB HL for the better ear or self-reported hearing impairment. Population based studies in English reported since 1970 with subjects aged 60 years and above were considered. 19 studies reporting audiometrically measured hearing loss and 5 studies reporting self-reported hearing impairment met our selection criteria. They did not allow establishing an integrative quantitative overview of prevalence rates over age and hearing loss for Europe, but they reflected the well-known patterns of a nonlinear increase of hearing loss with age affecting men more than women. Roughly 30% of men and 20% of women in Europe have a hearing loss of 30 dB HL or more at age 70 years, and 55% of men and 45% of women at age 80 years. The search confirms the known fact that ARHL is a major health concern in the aging population of Europe, but important information gaps became evident. They demonstrate the need of systematic and standardized collections of epidemiological data on hearing loss.

1. Introduction

The age distribution of the populations of developed countries has changed dramatically during the 20th century. The populations of Europe have become progressively older, particularly in the second half of the 20th century. A recent projection suggested that the majority of babies being born in this century will become 100 and more years old if the gain in longevity continues [1]. A steady linear growth of life expectancy to a undetermined biological limit was hypothesized in this work, but other studies considering different aspects and factors limiting duration of life [2] forecasted even a declining life expectancy due to factors such as the consequences of the increased prevalence of obesity [3]. Moreover, the prevalence of diseases such as cancer or dementia clearly increases because of longevity in our populations.

Fortunately, there is also good evidence that functional limitations and participation restrictions have diminished in the aged population along with the prolongation of life expectancy. Mobility and activities of daily living have improved. Age-related visual functional impairment has also decreased in the last decades, with cataract surgery contributing substantially to this improvement.

But what about hearing? Hearing loss (HL) is the most widespread sensory impairment in aging people. Hearing acuity declines with age - physiologically beginning by the third decade, predominately in the high frequencies. It begins to affect the frequencies of the speech spectrum within the fifth decade. Age-related hearing loss (ARHL) is primarily a hearing loss related to functional loss of sensory [4, 5] and neural elements [6], comparable to macular degeneration in the visual system rather than what is referred to as presbyopia, which is related to conductive disturbances of the optical system. Presbyopia can be corrected relatively easily by cataract surgery, whereas presbycusis exhibit multifaceted aspects involving also changes in neural structures including the central auditory processing. The cognitive and psychosocial consequences of HL are well described and well known [7], but population-based epidemiological knowledge or trends of the amount of HL and its health-related consequences are not well described.

Several aspects are of importance in this epidemiological context: differences in definition of HL, different measurements of HL, and difficulties in measuring the activity limitations and participation restrictions induced by HL. Trends of improvement of age-related HL either connected to longevity or by intervention have also not been identified clearly.

The aim of this review is to evaluate epidemiological data with respect to prevalence of ARHL in Europe and, if possible, to identify trends of prevalence over the last 50 years. Moreover, based on our findings and including additional aspects we will discuss procedures and strategies that may be included in future studies with the aim to improve epidemiologic knowledge of ARHL. Such knowledge may help to screen for ARHL, to induce earlier and more efficient interventions, and to reduce the burdens of ARHL.

2. Methods

Search methods: A systematic literature review was carried out with the specific aim to gain a picture of the epidemiology of ARHL in Europe.

Table 1 lists the sources and the search strategy. We searched the proposed core databases for a meta search in the category 'Medicine and Health'. These data bases partially overlap. Our search in the Library Catalogue of the University of Basel accessed the complete data base of the University of Basel (approximately 3 million titles) and those of 190 additional libraries of the Universities of Basel and Bern. The search was conducted in March and April 2010. Key words were entered in expressions with 'OR' when possible. Thesaurus function of PubMed (MeSH) was used to capture relevant alternative expressions for age related hearing loss.

Selection criteria: The results were further searched by hand. The references of relevant publications were systematically searched for additional relevant studies. We considered only population and epidemiological studies in English since 1970 with samples in European countries with subjects aged 60 years and above. For example, the study from Megighian et al. was excluded because of a non-representative proportion of 92% males [8]. A recent report from Davis et al. had to be excluded from table 4 because of the wide age group of 55-75 years [9].

Table 2 shows the two main standardized categorizations [10-12]. A recent report of the WHO proposed that disabling hearing impairment in adults should be defined as a permanent unaided hearing threshold level for the better ear of 41 dB or greater ($PTA_{0.5, 1, 2, 4} \geq 41$ dB HL) [13]. For reasons of homogeneity, we focus on prevalence data for the better ear (BEAR) and thresholds between 25 and 45 dB HL. Both audiometrically measured hearing loss (table 4) and self-reported hearing loss (table 5) were included. Some studies applied both methods of defining hearing loss.

3. Results

Numerical results of the search are summarized in **table 1**. The search terms offered 1150 results in Google scholar. These results were hand searched and compared to the hits in the library catalogues. In a second step, the references of the selected papers were searched for further relevant publications with no further results.

There is a paucity of European epidemiological studies on age related hearing loss in people aged 60 years and above [14, 15]. Apparent problems in comparing the available data are the heterogeneity of measures and cut-offs for grades of hearing impairment [14, 16].

Table 3 lists the 19 studies in alphabetic order that met our selection criteria.

We found two Italian studies in two different samples [17, 18], six Swedish studies in three different samples [19-24], two Norwegian studies in one sample [25, 26], four Finish studies in three samples [11, 12, 27, 28], two British studies in two samples [29, 30], one study in Sweden, Denmark and Finland with one of the aforementioned Finish samples [31], and a Dutch study [32].

Table 4 shows the results of another set of studies, again in alphabetical order. These studies describe self-reported hearing impairment. We found three British studies with three different samples [9, 29, 33], one study in Finland [27], one study in Denmark and Finland with the aforementioned Finish sample [31], one study in Sweden [34], and one study in Italy [35].

The study from Martini et al. is the only study that uses published hearing categories [36]. All other studies use either a single post-hoc dichotomized item to capture hearing impairments [27], a single post-hoc tripartite item, one binary item [33] , two binary items [9], or a single 5-point ordinal scale [29].

4. Discussion

Our systematic review of epidemiological data on prevalence of ARHL in Europe revealed more information gaps than information which would allow gaining a meaningful picture. Neither geographic distributions nor developments over time could be extracted to a reasonable degree. Most studies reported prevalence rates over age and frequency intervals; others reported data for single ages and frequencies. Because of this heterogeneity of the data we were unable to establish an integrative quantitative overview of prevalence rates over age

and hearing loss. Nevertheless, the studies do reflect the well-known patterns of a nonlinear increase of hearing loss with age affecting men more than women. If the data are crudely averaged and interpolated, roughly 30% of men and 20% of women in Europe have a hearing loss of 30 dB HL or more at age 70 years, and 55% of men and 45% of women at age 80 years.

These data confirm the known fact that ARHL is a major health concern in the aging population of Europe, but because of important information gaps it is difficult to devise any recommendations from them. They demonstrate the need of standardized collection of epidemiological data on hearing loss.

Difficulties in data evaluation - classification of HL

An evident and primary difficulty in comparing reported prevalence data are the different measures and cut-offs for hearing impairment [14, 16]. For example, the sample of the study in Denmark, Sweden and Finland consisted of 75-year-old subjects and results were subdivided in categories such as 'with minor difficulties' or 'with considerable difficulties' of hearing [31]. Other studies did not consistently report sample sizes [34], had small sample sizes [27], or summarized results over all age groups [35]. Many studies reported data with cut-offs of 25 dB and 35 dB HL. One study [17] proposes an adaptive threshold model that uses different thresholds as a function of frequency [37], but without applying these thresholds in the reported data.

Prevalence is relatively easy to establish in clearly defined states, such as whether a person is alive or dead, or if a pathohistological diagnosis of cancer exists. Hearing loss is a gradual condition and the limit between normal and not normal must be defined along a continuous scale. Obviously, the lack of a common definition prevents the availability of comparable prevalence data of HL.

For the sake of acceptable homogeneity and to obtain an overview, we focused on prevalence data for the better ear and a range of cut-offs for the definition of HL between 25 and 45 dB in the audiometric frequency range of 0.5 - 4 kHz. Tables 4 and 5 reveal that such limits cannot be regarded as a standard. In fact, it may be difficult to define any standard when evaluating the literature considered in this review. Even international classification systems such as those of the EU or the WHO differ considerably (table 2). Because any definition of HL has

unavoidably arbitrary elements, the acceptance and use of a standard definition of HL should be easy. It may be more the multitude of systems in use and the lack of knowledge of systems such as the WHO classification that hinders the regular use of them.

It seems to us that the use of the WHO classification in the collection of epidemiological data is a logical step which can help to improve such data in the future. Universal acceptance and propagation of the WHO classification system is recommended.

Difficulties in data evaluation - selection of populations

Similar arbitrary limits occur when defining the “age” for prevalence in ARHL. Again, tables 4 and 5 reveal that a standard age limit may be even more difficult to extract than the definition of HL. Even gender as a known significant influence on ARHL was not always clear. Only a portion of the studies explicitly reported prevalence rates for men and women, age groups, and hearing impairment categories [18, 20, 25, 27, 31]. Some studies either did not separate age and gender [11, 12, 17, 24, 28, 29, 32], did not calculate row and column prevalence for hearing impairments among other impairments [19], did not report sub-sample sizes for age groups [25], had overlapping age class limits [17], or had inconsistent hearing loss criteria between age groups [28]. Three studies did not report prevalence data even though it seemed possible judging from the description of the data [21-23, 26]. In the study of Smits et al. we derived approximate prevalence rates from a figure [32].

While international systems of HL classification exist, we are not aware of recommendations about the age limit defining ARHL. Even though the definition of such a limit may be biologically difficult if not impossible, it is highly desirable or even necessary to do so for epidemiological data. Again, for the sake of homogeneity this review sets a lower age limit of 60 years.

Connections between Aetiology and Epidemiology

Epidemiology and aetiology are intimately related in that population based endogenous factors will be exposed to geographically varying exogenous factors. One of our initial goals was to search for evidence linking epidemiology and aetiology.

As demonstrated in the previous sections, we have only a crude and patchy picture of the overall prevalence of ARHL in Europe. We know even less about the interaction of the

different aetiological factors leading to what is termed ARHL. A separate overview of the literature not reported here in details revealed that the most relevant factors in the aetiology of ARHL seem to be heredity (that may contribute up to 50%), noise, history of chronic middle ear inflammation, and cardiovascular factors including diabetes, smoking and hypertension; additional relevant factors are hormones (including gender), exposure to ototoxic medication or chemicals and co-morbidities. As an example, ARHL seems to be more prevalent in patients with rheumatologic disease [38, 39].

Given the vast gaps in epidemiological knowledge, it is not surprising that we were not able to establish meaningful connections between epidemiology and aetiological factors. Besides the complex biological background, it is also the lack of clinical methods able to differentiate and measure the contributions of the different pathophysiological mechanisms that limits our ability to gain meaningful data on epidemiology and aetiology. The differentiation between peripheral and central hearing disturbance is often difficult to assess [40, 41]. Patients with impairment of the central nervous system typically also exhibit deficits in other cognitive functions that may affect daily life additionally or more severely than hearing loss and unresolved questions of the reciprocal influence between hearing loss and cognition or isolation and central nervous function turn up.

Geographic distribution in prevalence

Most studies observing the demographic pattern of hearing loss were made in the northern and western part of Europe and there were only two studies in southern Europe [17, 18]. The findings of these two latter studies did not differ clearly from those of the other studies. Some additional data was found from Poland, the Netherlands, and Germany, but it was either inapplicable or not population based.

No geographic related pattern could be derived from these data. The relative absence of studies in the middle, southern and eastern parts of Europe may reflect more the differences in public health systems than attaching less importance to ARHL in these regions. Nevertheless, given the fact that vowels have a distinct perceptual advantage over consonants in determining speech intelligibility [42, 43], the different languages spoken in Europe may also influence epidemiological data of ARHL with respect to prevalence of self-reported hearing impairment.

Screening for ARHL?

Ideally, a comprehensive strategy to screen, assess, and improve ARHL would be based not only on crude prevalence data and projections of its longitudinal development, but on prevalence of aetiologically different subsets. It seems likely that ARHL with a primary aetiological background of vascular or metabolic factors will need different preventive measures than an ARHL primarily connected to factors such as noise or due to inflammatory middle ear disease. Given the biological variability of ARHL, it may be difficult to devise a universal approach for improving ARHL. Rehabilitation strategies may also have to consider the contribution of peripheral and central auditory aspects [41, 44]. Testing these components selectively is not well established and no experience of screening them exists.

Conclusion

This review of the literature did not allow gaining a clear picture of the prevalence of ARHL in Europe. Development of hearing loss over time and in conjunction with the increasing life expectancy is a major factor determining strategies of detection and correction of ARHL. The need for standardized procedures when collecting and reporting epidemiological data on HL has become evident. We recommend to use strictly the WHO classification of HL and to include regularly audiometric measure in population based health surveys.

References

1. Christensen, K., et al., *Ageing populations: the challenges ahead*. Lancet, 2009. **374**(9696): p. 1196-208.
2. Carnes, B., S. Olshansky, and D. Grahn, *Biological evidence for limits to the duration of life*. Biogerontology, 2003. **4**(1): p. 31-45.
3. Olshansky, S., et al., *A potential decline in life expectancy in the United States in the 21st century*. N Engl J Med, 2005. **352**(11): p. 1138-45.
4. Soucek, S., L. Michaels, and A. Frohlich, *Evidence for hair cell degeneration as the primary lesion in hearing loss of the elderly*. J Otolaryngol, 1986. **15**(3): p. 175-83.
5. Nelson, E. and R. Hinojosa, *Presbycusis: a human temporal bone study of individuals with downward sloping audiometric patterns of hearing loss and review of the literature*. Laryngoscope, 2006. **116**(9 Pt 3 Suppl 112): p. 1-12.
6. Frisina, R. and J. Walton, *Age-related structural and functional changes in the cochlear nucleus*. Hear Res, 2006. **216-217**: p. 216-23.
7. Monzani, D., et al., *Psychological profile and social behaviour of working adults with mild or moderate hearing loss*. Acta Otorhinolaryngol Ital, 2008. **28**(2): p. 61-6.
8. Megighian, D., et al., *Audiometric and epidemiological analysis of elderly in the Veneto region*. Gerontology, 2000. **46**(4): p. 199-204.
9. Davis, A., et al., *Acceptability, benefit and costs of early screening for hearing disability: a study of potential screening tests and models*. Health Technol Assess, 2007. **11**(42): p. 1-294.
10. Martini, A., et al., *Audiometric patterns of genetic non-syndromal sensorineural hearing loss*. Audiology, 1997. **36**(4): p. 228-36.
11. Uimonen, S., et al., *Do we know the real need for hearing rehabilitation at the population level? Hearing impairments in the 5- to 75-year-old cross-sectional Finnish population*. Br J Audiol, 1999. **33**(1): p. 53-9.
12. Uimonen, S., et al., *Hearing in 55 to 75 year old people in northern Finland--a comparison of two classifications of hearing impairment*. Acta Otolaryngol Suppl, 1997. **529**: p. 69-70.
13. World Health Organization and Regional Office for South-East Asia, *Deafness and Hearing Impairment Survey - Report of the Consultative Meeting of Principal Investigators SEARO, New Delhi, 7- 9 May 2001*. 2001, WHO: New Delhi.
14. Cruickshanks, K.J., W. Zhan, and w. Zhong, *Epidemiology of age-related hearing impairment*, in *The Aging Auditory System*, S. Gordon-Salant, et al., Editors. 2010, Springer Science+Business Media. p. 259-274.
15. Davis, A., K. Davis, and P. Smith, *The prevalence of deafness and hearing impairment*, in *Ballantyne's Deafness*, J. Graham and M. Martin, Editors. 2009, Wiley. p. 6-19.
16. Jerger, J., et al., *Hearing impairment in older adults: new concepts*. J Am Geriatr Soc, 1995. **43**(8): p. 928-35.
17. Bedin, E., et al., *Age-related hearing loss in four Italian genetic isolates: an epidemiological study*. Int J Audiol, 2009. **48**(7): p. 465-72.
18. Quaranta, A., G. Assennato, and V. Sallustio, *Epidemiology of hearing problems among adults in Italy*. Scand Audiol Suppl, 1996. **42**: p. 9-13.
19. Bergmann, B. and U. Rosenhall, *Vision and hearing in old age*. Scandinavian Audiology, 2001. **30**(4): p. 255-263.
20. Johansson, M.S. and S.D. Arlinger, *Prevalence of hearing impairment in a population in Sweden*. Int J Audiol, 2003. **42**(1): p. 18-28.

21. Jonsson, R. and U. Rosenhall, *Hearing in advanced age. A study of presbycusis in 85-, 88- and 90-year-old people*. Audiology, 1998. **37**(4): p. 207-18.
22. Jonsson, R., et al., *Auditory function in 70- and 75-year-olds of four age cohorts. A cross-sectional and time-lag study of presbycusis*. Scand Audiol, 1998. **27**(2): p. 81-93.
23. Moller, M.B., *Hearing in 70 and 75 year old people: results from a cross sectional and longitudinal population study*. Am J Otolaryngol, 1981. **2**(1): p. 22-9.
24. Rosenhall, U. and A.K. Karlsson Espmark, *Hearing aid rehabilitation: what do older people want, and what does the audiogram tell?* Int J Audiol, 2003. **42 Suppl 2**: p. 2S53-7.
25. Borchgrevink, H.M., K. Tambs, and H.J. Hoffman, *The Nord-Trondelag Norway Audiometric Survey 1996-98: unscreened thresholds and prevalence of hearing impairment for adults > 20 years*. Noise Health, 2005. **7**(28): p. 1-15.
26. Engdahl, B., et al., *Screened and unscreened hearing threshold levels for the adult population: results from the Nord-Trondelag Hearing Loss Study*. Int J Audiol, 2005. **44**(4): p. 213-30.
27. Hietanen, A., et al., *Changes in hearing in 80-year-old people: a 10-year follow-up study*. Int J Audiol, 2004. **43**(3): p. 126-35.
28. Rahko, T., et al., *Prevalence of handicapping hearing loss in an aging population*. Ann Otol Rhinol Laryngol, 1985. **94**(2 Pt 1): p. 140-4.
29. Davis, A.C., *The prevalence of hearing impairment and reported hearing disability among adults in Great Britain*. Int J Epidemiol, 1989. **18**(4): p. 911-7.
30. Wilson, P.S., D.M. Fleming, and I. Donaldson, *Prevalence of hearing loss among people aged 65 years and over: screening and hearing aid provision*. Br J Gen Pract, 1993. **43**(375): p. 406-9.
31. Hietanen, A., et al., *Hearing among 75-year-old people in three Nordic localities: a comparative study*. Int J Audiol, 2005. **44**(9): p. 500-8.
32. Smits, C. and T. Houtgast, *Results from the Dutch speech-in-noise screening test by telephone*. Ear Hear, 2005. **26**(1): p. 89-95.
33. Chou, K.L., *Combined effect of vision and hearing impairment on depression in older adults: evidence from the English Longitudinal Study of Ageing*. J Affect Disord, 2008. **106**(1-2): p. 191-6.
34. Rosenhall, U., R. Jonsson, and O. Soderlind, *Self-assessed hearing problems in Sweden: a demographic study*. Audiology, 1999. **38**(6): p. 328-34.
35. Martini, A., et al., *Hearing in the elderly: a population study*. Audiology, 2001. **40**(6): p. 285-93.
36. Sanders, D.A., *Hearing aid orientation and counselling*, in *Amplification for the hearing-impaired*, M.C. Pollack, Editor. 1975, Grune & Stratton: New York. p. 121-128.
37. Sakamoto, M., et al., *Average thresholds in the 8 to 20 kHz range as a function of age*. Scand Audiol, 1998. **27**(3): p. 189-92.
38. Murdin, L., et al., *Hearing difficulties are common in patients with rheumatoid arthritis*. Clin Rheumatol, 2008. **27**(5): p. 637-40.
39. George, D. and S. Pradhan, *Idiopathic sensorineural hearing disorders in adults--a pragmatic approach*. Nat Rev Rheumatol, 2009. **5**(9): p. 505-12.
40. Gates, G. and G. Popelka, *Neural presbycusis: a diagnostic dilemma*. Am J Otol, 1992. **13**(4): p. 313-7.
41. Otto, W. and G. McCandless, *Aging and auditory site of lesion*. Ear Hear, 1982. **3**(3): p. 110-7.
42. Kewley-Port, D., T. Burkle, and J. Lee, *Contribution of consonant versus vowel*

- information to sentence intelligibility for young normal-hearing and elderly hearing-impaired listeners. J Acoust Soc Am, 2007. 122(4): p. 2365-75.*
43. Fogerty, D. and D. Kewley-Port, *Perceptual contributions of the consonant-vowel boundary to sentence intelligibility. J Acoust Soc Am, 2009. 126(2): p. 847-57.*
 44. Gates, G. and J. Mills, *Presbycusis. Lancet, 2005. 366(9491): p. 1111-20.*
 45. Davis, A., *Hearing in adults. 1995, London: Whurr.*

Table 1 Search Strategy

Database¹	search term	number of results	decision	hand search in results
“Google Scholar” via VPN ² of University of Basel	prevalence AND ("hearing loss")	<i>in document:</i> 31'900 <i>in title:</i> 217	restricted search in results for "in title"	4 relevant
	epidemiology AND ("hearing loss")	<i>in document:</i> 21'400 <i>in title:</i> 59	restricted search in results for "in title"	0 additional results
	(prevalence OR epidemiology OR epidemiological) AND ("age related hearing loss" OR "presbiacosis") ³	<i>in document:</i> 1150 <i>in title:</i> 5	hand search in all results	38 additional results
PubMed	see above	<i>in document:</i> 59	hand search in all results	0 additional results
Recherche-Portal provided by Hauptbibliothek Universität Zürich ⁴	see above	<i>in document:</i> 690	hand search in all results	10 additional results
Eurostat; domain Health (public health/safety at work) ⁵	all above mentioned search term components	no results		

Note:

¹ Search in databases follows a chronological order

² Virtual Private Network, allowed direct access to hosting organizations such as libraries and subscribed journals

³ combinations of search terms allows more results, ARHL (Age Related Hearing Loss) and presbiacosis were chosen as most appropriate target expression

⁴ Meta Search in "Medicine and Health"/ “Core Databases”: CINAHL (EBSCO), Cochrane Library (Wiley), EMBASE.com, MEDLINE (OvidSP), MEDLINE incl. Premedline, Premedline (OvidSP), PsycINFO (OvidSP), PubMed / Medline, Springer Protocols, Web of Science (ISI) XML Gateway

⁵ Search term combinations not supported

Table 2 Standardized hearing loss categories

Categorization	(1) 'EU' classification	(2) 'WHO' classification
normal	$\text{dBHL} < 21$	$\text{dBHL} < 26$
mild	$21 \leq \text{dBHL} < 39$	$26 \leq \text{dBHL} < 40$
moderate	$40 \leq \text{dBHL} < 69$	$41 \leq \text{dBHL} < 60^{\text{a}}$
severe	$70 \leq \text{dBHL} < 94$	$61 \leq \text{dBHL} < 80$
profound	$94 < \text{dBHL}$	$80 < \text{dBHL}$
Note: ^a according to the WHO, hearing impairment for the better ear of 41 dB or above have been defined as disabling		

Table 3 Measured prevalence of age-related hearing loss ≥ 30 dBHL in elderly 60+ years: study overview

Reference	Study	Prevalence with nearest cut-offs for HL and age ^a
Bedin et al, 2009; [17]	Italy; multidisciplinary project in several genetically isolated villages	Inapplicable data
Bergmann & Rosenhall, 2001; [19]	See Jonsson & Rosenhall, 1998; Jonsson et al., 1998; [21, 22]	> 19% (70y at 30-39dB)
Borchgrevink et al., 2005; [25]	Norway; Supplement to the Nord-Trøndelag Health Survey (Helseundersøkelsen i Nord-Trøndelag, HUNT II); 1996-1998	> 14.2% (60-64y at 35dB)
Davis, 1989; [29]	UK; national epidemiological study; MRC Institute of Hearing Research); 1980-1986	> 7.4% (61-70y at 45dB)
Davis, 1995; [45]	see Davis (1989)	> 24.5% (61-70y at 30dB)
Engdahl et al., 2005; [26]	Norway; Regular part of the Nord-Trøndelag Health Survey (Helseundersøkelsen i Nord-Trøndelag, HUNT II); 1995-1997	Inapplicable data ^b
Hietanen et al., 2005; [31]	Denmark, Sweden, Finland; NORA (Nordic Research on Aging); 1989-1991	>16.5% (75y at 40-69dB)
Hietanen et al., 2004; [27]	Finland; 1990-2000	>28.3% (80y at 40-69dB)
Johansson & Arlinger, 2003; [20]	Sweden; 1998	> 8.8% (60-70y at 35dB)
Jonsson & Rosenhall, 1998; Jonsson et al., 1998; [21, 22]	Sweden; gerontological and geriatric population study (H70); Subjects born in period 1st July 1901 - 30 June 1902; 1971-1992	Inapplicable data ^c

Moller, 1981; [23]	Sweden; gerontological and geriatric population study (H70 - H75); 1971-1992; also see Jonsson & Rosenhall, 1998 [21]	9% (70y at 35dB)
Quaranta et al., 1996; [18]	Italy;	6.7% (61-70y at 45dB)
Rahko et al., 1985; [28]	Finland; national "Project 65" study; 1975-1978	10.3% (65y at 30dB)
Rosenhall et al., 2003; [24]	Sweden; N=237 "Q/H70"; 1992-1993; N=168 "NORA"; 1991-1992; N=154 "Gothenburg"; 1991-1990	24% (70y at 30-39dB)
Smits & Houtgast, 2005; [32]	The Netherlands; "National Hearing" test (automatic speech-in-noise screening test by telephone); 2005	Inapplicable data ^d
Uimonen et al., 1999; [11]	Finland; Hearing loss classification study	Inapplicable data ^e
Uimonen et al., 1997; [12]	See Uimonen et al., 1999; [11]	Inapplicable data ^f
Wilson et al., 1993; [30]	UK; 1989-1990	54.3% (≥65y at 35dB)

Note:

^a minimum prevalence in men and women for minimum better ear hearing loss of 30dB and lower age interval border of 60 years

^b graphical data report with threshold curves for separate pure tone frequencies

^c data report quartiles for cohorts and pure tone frequencies

^d data report speech reception thresholds in noise

^e hearing loss interval (26-40dB) falls below target cut-off (30dB)

^f age interval (55-75 years) falls below target cut-off (60 years)

Table 4 Self-reported prevalence of age-related hearing loss in elderly 60+ years: study overview

Reference	Study	Prevalence: impairments of any kind ^a
Chou, 2008; [33]	UK; English Longitudinal Study of Ageing (ELSA), wave I (2002-2003) and II (2005)	> 18.9% (≥65 years)
Davis et al., 2007; [9]	UK; National Study of Hearing (NSH)	>40.3% (≥75 years)
Davis, 1989; [29]	See table 4	> 14.6% (> 60 years)
Hietanen et al, 2005; [31]	Denmark, Sweden, Finland; NORA (Nordic Research on Aging); 1989-1991	> 27.7% (75 years)
Hietanen et al., 2004; [27]	Finland; 1990-2000	> 44.7% (80 years)
Martini et al., 2001; [35]	Italy; 1989	> 8.1% (>60 years)
Rosenthal et al., 1999; [34]	Sweden; Part of the Swedish Survey of Living Conditions (ULF); conducted by Statistics Sweden (SCB); 1986-1993	> 16% (>64 years)

Note:

^a minimum prevalence in men and women for subjective hearing loss in study-specific impairment criterion

Table 1 Search Strategy

Database¹	search term	number of results	decision	hand search in results
“Google Scholar” via VPN ² of University of Basel	prevalence AND ("hearing loss")	<i>in document:</i> 31'900 <i>in title:</i> 217	restricted search in results for "in title"	4 relevant
	epidemiology AND ("hearing loss")	<i>in document:</i> 21'400 <i>in title:</i> 59	restricted search in results for "in title"	0 additional results
	(prevalence OR epidemiology OR epidemiological) AND ("age related hearing loss" OR "presbiacusis") ³	<i>in document:</i> 1150 <i>in title:</i> 5	hand search in all results	38 additional results
PubMed	see above	<i>in document:</i> 59	hand search in all results	0 additional results
Recherche-Portal provided by Hauptbibliothek Universität Zürich ⁴	see above	<i>in document:</i> 690	hand search in all results	10 additional results
Eurostat; domain Health (public health/safety at work) ⁵	all above mentioned search term components	no results		

Note:

¹ Search in databases follows a chronological order

² Virtual Private Network, allowed direct access to hosting organizations such as libraries and subscribed journals

³ combinations of search terms allows more results, ARHL (Age Related Hearing Loss) and presbiacusis were chosen as most appropriate target expression

⁴ Meta Search in "Medicine and Health"/ "Core Databases": CINAHL (EBSCO), Cochrane Library (Wiley), EMBASE.com, MEDLINE (OvidSP), MEDLINE incl. Premedline, Premedline (OvidSP), PsycINFO (OvidSP), PubMed / Medline, Springer Protocols, Web of Science (ISI) XML Gateway

⁵ Search term combinations not supported

Table 2 Standardized hearing loss categories

Categorization	(1) 'EU' classification	(2) 'WHO' classification
normal	dBHL < 21	dBHL < 26
mild	$21 \leq \text{dBHL} < 39$	$26 \leq \text{dBHL} < 40$
moderate	$40 \leq \text{dBHL} < 69$	$41 \leq \text{dBHL} < 60^a$
severe	$70 \leq \text{dBHL} < 94$	$61 \leq \text{dBHL} < 80$
profound	$94 < \text{dBHL}$	$80 < \text{dBHL}$
Note: ^a according to the WHO, hearing impairment for the better ear of 41 dB or above have been defined as disabling		

Table 3 Measured prevalence of age-related hearing loss ≥ 30 dBHL in elderly 60+ years: study overview

Reference	Study	Prevalence with nearest cut-offs for HL and age ^a
Bedin et al, 2009; [17]	Italy; multidisciplinary project in several genetically isolated villages	Inapplicable data
Bergmann & Rosenhall, 2001; [19]	See Jonsson & Rosenhall, 1998; Jonsson et al., 1998; [21, 22]	> 19% (70y at 30-39dB)
Borchgrevink et al., 2005; [25]	Norway; Supplement to the Nord-Trøndelag Health Survey (Helseundersøkelsen i Nord-Trøndelag, HUNT II); 1996-1998	> 14.2% (60-64y at 35dB)
Davis, 1989; [29]	UK; national epidemiological study; MRC Institute of Hearing Research); 1980-1986	> 7.4% (61-70y at 45dB)
Davis, 1995; [45]	see Davis (1989)	> 24.5% (61-70y at 30dB)
Engdahl et al., 2005; [26]	Norway; Regular part of the Nord-Trøndelag Health Survey (Helseundersøkelsen i Nord-Trøndelag, HUNT II); 1995-1997	Inapplicable data ^b
Hietanen et al., 2005; [31]	Denmark, Sweden, Finland; NORA (Nordic Research on Aging); 1989-1991	>16.5% (75y at 40-69dB)
Hietanen et al., 2004; [27]	Finland; 1990-2000	>28.3% (80y at 40-69dB)
Johansson & Arlinger, 2003; [20]	Sweden; 1998	> 8.8% (60-70y at 35dB)
Jonsson & Rosenhall, 1998; Jonsson et al., 1998; [21, 22]	Sweden; gerontological and geriatric population study (H70); Subjects born in period 1st July 1901 - 30 June 1902; 1971-1992	Inapplicable data ^c

Moller, 1981; [23]	Sweden;gerontological and geriatric population study (H70 - H75); 1971-1992; also see Jonsson & Rosenhall, 1998 [21]	9% (70y at 35dB)
Quaranta et al., 1996; [18]	Italy;	6.7% (61-70y at 45dB)
Rahko et al., 1985; [28]	Finland; national "Project 65" study; 1975-1978	10.3% (65y at 30dB)
Rosenhall et al., 2003; [24]	Sweden; N=237 "Q/H70";1992-1993; N=168 "NORA"; 1991-1992; N=154 "Gothenburg"; 1991-1990	24% (70y at 30-39dB)
Smits & Houtgast, 2005; [32]	The Netherlands; “National Hearing” test (automatic speech-in-noise screening test by telephone); 2005	Inapplicable data ^d
Uimonen et al., 1999; [11]	Finland; Hearing loss classification study	Inapplicable data ^e
Uimonen et al., 1997; [12]	See Uimonen et al., 1999; [11]	Inapplicable data ^f
Wilson et al., 1993; [30]	UK; 1989-1990	54.3% (≥65y at 35dB)

Note:

^a minimum prevalence in men and women for minimum better ear hearing loss of 30dB an lower age interval border of 60 years

^b graphical data report with threshold curves for separate pure tone frequencies

^c data report quartiles for cohorts and pure tone frequencies

^d data report speech reception thresholds in noise

^e hearing loss interval (26-40dB) falls below target cut-off (30dB)

^f age interval (55-75 years) falls below target cut-off (60 years)

Table 4 Self-reported prevalence of age-related hearing loss in elderly
60+ years: study overview

Reference	Study	Prevalence: impairments of any kind ^a
Chou, 2008; [33]	UK; English Longitudinal Study of Ageing (ELSA), wave I (2002-2003) and II (2005)	> 18.9% (≥65 years)
Davis et al., 2007; [9]	UK; National Study of Hearing (NSH)	>40.3% (≥75 years)
Davis, 1989; [29]	See table 4	> 14.6% (> 60 years)
Hietanen et al, 2005; [31]	Denmark, Sweden, Finland; NORA (Nordic Research on Aging); 1989-1991	> 27.7% (75 years)
Hietanen et al., 2004; [27]	Finland; 1990-2000	> 44.7% (80 years)
Martini et al., 2001; [35]	Italy; 1989	> 8.1% (>60 years)
Rosenthal et al., 1999; [34]	Sweden; Part of the Swedish Survey of Living Conditions (ULF); conducted by Statistics Sweden (SCB); 1986-1993	> 16% (>64 years)

Note:

^a minimum prevalence in men and women for subjective hearing loss in study-specific impairment criterion

Prevalence of age-related hearing loss (ARHL) in Europe – A Review

*Thomas Roth¹, Dirk Hanebuth^{2,3,4}, Rudolf Probst¹

¹ University Hospital of Zurich, Department of Otorhinolaryngology, Head and Neck Surgery, Frauenklinikstr. 24, 8091 Zürich, Switzerland

² Swiss Tropical and Public Health Institute, Department Epidemiology and Public Health, Unit Gender and Health, Socinstrasse 57, 4002 Basel, Switzerland

³ University of Basel, Petersplatz 1, 4003 Basel, Switzerland

⁴ ETH Zurich, Department of Computer Science, Universitätstrasse 6, 8092 Zürich

*Corresponding author: thomas.roth@usz.ch

Acknowledgment:

- This study was supported by the “Assessment of Hearing in Elderly: Aging and Degeneration” (AHEAD): AHEAD III was a project with several aims concerning age-related hearing loss. For further information see the weblink at www.ahead.polimi.it.
- We thank Fran Harris as a native English speaker for her linguistic corrections and suggestions.

Parts of this review were presented at the International Conference on Adult Hearing Screening in Cernobbio, Italy (June 10-12, 2010).

Keywords:

Presbycusis, age-related hearing loss, auditory system, cochlear function, epidemiology, Europe, AHEAD